



Provider Checklist-Outpatient – Cystoscopy/Cystourethroscopy

(CPT Code: 52000, 52005, 52204, 52214, 52224, 52234, 52240, 52260, 52265, 52276, 52281, 52285, 52310, 52315, 52320, 52332)

All Indications [*One has to be present]

*Hematuria [**One has to be present**]

Gross hematuria [All have to be present]

Blood by urine dipstick Urine culture negative Imaging study nondiagnostic for etiology of hematuria

Painless microscopic hematuria [All have to be present]

RBCs ≥ 3/HPF(High Power Field) No RBC casts by U/A UTI excluded [**One has to be present**]

> -1 Urine dipstick negative for nitrite and leukocyte esterase -2 Urine culture negative

Imaging study nondiagnostic for etiology of hematuria

*Obstructive symptoms [All have to be present]

Male patient ≤ 50 Symptoms [**One has to be present**]

> Decreased size/force of urine stream Hesitancy Intermittency Sense of post voiding fullness Straining on urination

Prostate exam [Both have to be present]

Nontender/not boggy by DRE (Digital Rectal Exam) No nodule/induration by DRE

Urine culture negative





Continued symptoms after Rx [**One has to be present**]

 $Abx \ge 3$ wks Alpha-blocker ≥ 3 wks

*Irritative symptoms w/o infection [One has to be present]

Males [All have to be present]

Symptoms [Both have to be present]

-1 Irritative symptoms [One has to be present]

A) UrgencyB) FrequencyC) DysuriaD) Urge incontinence

-2 No obstructive symptoms [All have to be present]

- A) No decreased size/force of urine stream
- B) No hesitancy
- C) No intermittency
- D) No sense of post voiding fullness
- E) No straining on urination

Prostate nontender/not boggy by DRE

Urine culture negative

Urodynamic evaluation normal

Females [All have to be present]

Symptoms [**One has to be present**]

- -1 Urgency \ge 4 wks
- -2 Frequency \ge 4 wks
- -3 Dysuria \ge 4 wks
- -4 Urge incontinence \geq 4 wks





Findings [Both have to be present]

-1 Urethral opening normal -2 No vaginitis

Urine culture negative

*Recurrent cystitis/pyelonephritis by culture [One has to be present]

Males [One has to be present]

Cystitis $\ge 2x \text{ w/in } 1 \text{ yr}$ Pyelonephritis ≥ 1 episode

Females [One has to be present]

Cystitis $\ge 4x$ w/in 1 yr Cystitis on Abx suppressant Rx Pyelonephritis ≥ 2 episodes

*Ureteroscopic lithotripsy for ureteral stone ≥ 5 mm [**One has to be present**]

Proximal ureter/renal pelvis Distal ureter [**One has to be present**]

> Complete obstruction with hydronephrosis by imaging Continued partial obstruction by imaging after adequate hydration ≥ 4 wks

Continued pain/vomiting after Rx > 24 hrs [**Both have to be present**]

-1 Hydration-2 Medication [One has to be present]

A) AnalgesicsB) Antiemetics

UTI by culture

*Basket retrieval of ureteral stone ≥ 5 mm [**Both have to be present**]

Distal ureter Persistent Sx/findings [**One has to be present**]





Complete obstruction with hydronephrosis by imaging Continued partial obstruction by imaging after adequate hydration ≥ 4 wks

Continued pain/vomiting after Rx > 24 hrs [**Both have to be present**]

-1 Hydration-2 Medication [One has to be present]

A) AnalgesicsB) Antiemetics

UTI by culture

*Placement of ureteral catheter/stent [One has to be present]

Ureteral obstruction by CT/US/IVP Ureteral stricture [**One has to be present**]

> Hydronephrosis by CT/US/IVP Pyelonephritis ≥ 2x w/in 1 yr Prior to lithotripsy

Prior to lithotripsy with stone > 1.5 cm

*Superficial bladder cancer [One has to be present]

TURBT (Transurethral Resection Bladder Tumor) planned Patient too ill/refuses cystectomy

- *Periodic assessment with treated bladder cancer
- *Positive urine cytology
- *Preoperative evaluation for ureteral reimplantation
- *Intravesical foreign body removal