

### cFluor™ Immunoprofiling Kit

Technical Data Sheet

# Cytek cFluor™ Immunoprofiling Kit

### **PRODUCT DETAILS**

Catalog number: R7-40000 (25 Tests)

Category: Immunophenotyping

Format: cFluor conjugated antibodies in individual tubes

| Target       | Clone   | Fluorochrome               | Item number |
|--------------|---------|----------------------------|-------------|
| CD3          | SK7     | cFluor V420                | R7-20054    |
| CD4          | SK3     | cFluor R780 <sup>1</sup>   | R7-20084    |
| CD8          | SK1     | cFluor B515 <sup>4</sup>   | R7-20036    |
| CD14         | M5E2    | cFluor V450                | R7-20004    |
| CD16         | 3G8     | cFluor R668 <sup>4</sup>   | R7-20070    |
| CD19         | HIB19   | cFluor BYG710              | R7-20010    |
| CD25         | BC96    | cFluor BYG781 <sup>3</sup> | R7-20080    |
| CD27         | QA17A18 | cFluor R840 <sup>2</sup>   | R7-20082    |
| CD45         | HI30    | cFluor V547 <sup>4</sup>   | R7-20012    |
| CD45RA       | HI100   | cFluor B690                | R7-20086    |
| CD56         | 5.1H11  | cFluor R720 <sup>4</sup>   | R7-20090    |
| CD127        | A019D5  | cFluor R659 <sup>3</sup>   | R7-20078    |
| CD197 (CCR7) | G043H7  | cFluor BYG575 <sup>3</sup> | R7-20076    |
| IgD          | IA6-2   | cFluor BYG667 <sup>3</sup> | R7-20088    |

| Test Dilution: | 5µl per test  |
|----------------|---|
| Application:   | Flow cytometry  |
| Formulation:   | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% BSA |
| Storage:       | 2-8°C and protected from light.   |
| _              | Do not freeze   |

#### **PRODUCT DESCRIPTION**

Cytek cFluor Immunoprofiling Kit allow for the identification of helper T cells, cytotoxic T cells, B cells, NK cells and monocytes in human peripheral blood mononuclear cells and in whole blood. The reagents in this kit help to distinguish different subsets of T, B and NK cells; including regulatory T cells, naïve T cells, activated T cells, memory T cells, effector T cells, naïve B cell, memory B cells, nonclassical and classical monocytes.

CD3 is expressed on all mature T cells, NK T cells, and some thymocytes. CD3, a part of the CD3/T cell receptor complex, plays a role in antigen recognition, signal transduction, and T cell activation.

CD4, is expressed on most thymocytes, a major subset of T cells, and on monocytes/macrophages. Functionally, CD4 is associated with thymic differentiation, in conjunction with MHC class II molecules in antigen recognition and with signal transduction.



## cFluor™ Immunoprofiling Kit

### Technical Data Sheet

CD8 is found on thymocytes, on a subset of T cells and on NK cells. This molecule acts as a co-receptor in MHC class I molecules in antigen recognition, has a role in T cell activation and in thymic differentiation.

CD14 is also known as a high affinity LPS receptor and is highly expressed on monocytes and macrophages. It is also expressed on granulocytes, but at a lower level. In addition, CD14 is found on interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells.

CD16 is expressed on NK cells, monocytes and macrophages in the form of CD16a. Another form of CD16, CD16b is expressed on neutrophils. CD16 engagement of IgG leads to NK cell activation, antibody-dependent cell-mediated cytotoxicity (ADCC) and phagocytosis

CD19 is expressed in the B cell lineage, from pro-B to blastoid B cells. However, it is absent on plasma cells. It is also expressed on follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation.

CD25 is a low affinity IL-2 receptor that is expressed on progenitor lymphocytes, activated T & B cells, and monocytes, as well as on regulatory T cells. When CD25 is in association with CD122, IL-2 receptor beta chain and with CD132, the common gamma chain, the resulting complex is the high-affinity IL-2 receptor. IL-2 receptor signaling is key lymphocyte proliferation and survival.

CD27 is a lymphocyte specific member of the TNF receptor superfamily. CD27 is expressed on a subset of thymocytes, mature T cells. The expression further upregulated in T cell activation. Subpopulations of B cells and NK also express CD27. In T cell-B cell interaction, CD27 binds to CD70 to provide costimulation to result in T cell activation and B cell differentiation and proliferation.

CD45 is expressed on all hematopoietic cells, except erythrocytes and platelets. CD45 is a signaling molecule that is involved in cellular proliferation, differentiation and in regulation of immune cell functions.

CD45RA is an isoform of CD45 due to alternate splicing. CD45RA is expressed on naïve or resting CD4+ and CD8+ T cells, as well as on B cells and monocytes.

CD56, within the hematopoietic system is expressed on NK cells and NKT cells, a subset of T cells. In the nervous system, CD56 is expressed by neurons and plays a role in the homotypic adhesion of neural cells.

CD127, when it heterodimerizes with the common gamma chain (CD132) forms the IL-7 receptor. This receptor is involved in the development of B cells in the bone marrow, and the control of T cell proliferation in the periphery. CD127 expression is downregulated in regulatory T cells.

CD197 (CCR7) is a chemokine receptor for chemokines CCL19 and CCL21. CCR7 and its ligands are involved in trafficking and migration of immune cells into immunological milieus. For example, B cell migration and organization into secondary lymphoid tissues, and migration of memory T cells into inflamed tissue. CCR7 is expressed on T cells and the detection of CCR7, in combination with CD45RA, can be used to distinguish naive T cells from central and effector memory T cells.

IgD is a molecule that is expressed on naive B cells. IgD is expressed on B cells upon exiting the bone marrow to the periphery.



### cFluor™ Immunoprofiling Kit

Technical Data Sheet

#### **REFERENCES**

Clevers H et al. Ann Rev Immunol. 1988 **6**, 629 (1988) Evans RL et al. Immunol. **78**, 544 (1981) Cole DK et al. Cell Mol Immunol. **1**, 81 (2004) Ferrero E et al. J. Immunol. **145**, 331 (1990) Wirthmueller U et al. J Exp Med. **175**, (1992) Dennis O et al. J Immunol. **170**, 73 (2003) Shevach E. Nat Rev Immunol. **2**, 389 (2002) Vossen M et al. J Immunol. **180**, 3739 (2008) Fritsch R et al. J Immunol. 175, 6489 (2005) Loken MR et al. Cytometry. **11**, 453 (1990) Kumar, Connors & Farber. Immunity. **48**, 202 (2018) Poli A et al. Immunology. **126**, 458 (2009)

Carrette, F & Surh, C. Semin Immunol. **24,** 209 (2012) Mahnke Y et al. Eur J Immunol. **43,** 2797 (2013)

#### **RECOMMENDED USAGE**

Goodwin R et al. Cell. **73**, 447 (1993)

Human peripheral mononuclear cells (PBMC) or whole blood have been tested to validate the performance of this kit. For staining procedures, product data, and gating strategy, please refer to the Reagents and Protocols sections of our website at www.cytekbio.com.

For Research Use Only. Not intended for use in diagnostic procedures.

 $<sup>^{1}</sup>$  Cytek dye cFluor R780 and its conjugates were developed, manufactured and commercialized by BioLegend, Inc. under the trademark APC/Fire750<sup>™</sup>

<sup>&</sup>lt;sup>2</sup> Cytek dye cFluor R840 and its conjugates were developed, manufactured and commercialized by BioLegend, Inc. under the trademark APC/Fire810™

<sup>&</sup>lt;sup>3</sup> Fluor conjugated antibody manufactured and supplied by BioLegend Inc.

<sup>&</sup>lt;sup>4</sup>cFluor™ V547, cFluor™ B515, cFluor™ B532, cFluor™ R668 and cFluor™ R720 are equivalent to CF®405L, CF®488A, CF®503, CF®647 and CF®700 respectively, manufactured and provided by Biotium, Inc. under an Agreement between Biotium and Cytek (LICENSEE). The manufacture, use, sale, offer for sale, or import of the product is covered by one or more of the patents or pending applications owned or licensed by Biotium. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. No right under any other patent claim, no right to perform any patented method, and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is conveyed expressly, by implication, or by estoppel.